

DESIGN OF DERMA STICKS OF ABRUS PRECATORIUS FOR THE TREATMENT OF GYNECOLOGICAL SKIN INFECTIONS

**K. Purushotham Rao¹, Savita Sonawane², P. Sagare³, Pratima S⁴, Anuradha Patil³,
Rajsree Ingin⁵**

¹HKEs College of Pharmacy, MR Medical College, Gulbarga. K. S.

²College of Pharmacy, Solapur University, Solapur., M. S.

³MR Medical College and General Hospital-Gulbarga-K. S.

⁴KBN Medical College and General Hospital, Gulbarga. K. S.

⁵Govt Medical College and General Hospital, Gulbarga., K. S.

Article Received on
04 Oct. 2016,

Revised on 24 Oct. 2016,
Accepted on 15 Nov. 2016

DOI: 10.20959/wjpr201612-7442

***Corresponding Author**

K. Purushotham Rao

HKEs College of Pharmacy,
MR Medical College,
Gulbarga – 585105.

ABSTRACT

Candidosis (Candidiasis, moniliasis) is an infection of the skin, mucosa and rarely of the internal organs, caused by yeast like fungus *Candida albicans*, normally present in vagina. *Candida albicans* is an ovoid or spherical budding cell, which produces pseudo-mycelia both in culture and in tissues. *Candida* species are normal inhabitants of the skin and mucosa. Over the last two decades, there has been a dramatic increase in the rate of superficial and invasive fungal infections. Approximately three-quarters of all women experience at least one episode of vulva-vaginal candidiasis during their life time and nearly half of them suffer

from multiple episode. The manifestations of vulva vaginal candidiasis are often painful and uncomfortable and can include intense itching, irritation, vaginal discharge and dysuria. Gynecological skin disorders referring to inflammatory and infectious conditions affecting the vaginal mucosa and vulvitis often accompanies vaginal pain, itching and burning sensation. The topical drug delivery systems available for the treatment have several disadvantages like greasiness, inconvenient to store and requires applicator or use of fingertip, which may lead to contamination. Therefore, it was found essential to find an alternative to counter all the above disadvantages effectively and hence in the present work, formulation and development of medicated sticks has been planned with the herbal *Abrus Precatorius* drug which is very well known for the anti bacterial and anti fungal activity. The preparation and characterization of medicated sticks was carried out in four phases. Phase I

studies includes preparation of medicated sticks using the ointment bases with varied concentrations of waxes. Phase II studies involves characterization of prepared medicated derma sticks like weight, thickness and length. Phase III studies involves stability studies of prepared formulations. Phase IV Studies involves anti microbial studies by zone inhibition method. Phase V: Primary skin irritation studies carried out on rabbits and guinea pigs and in healthy human volunteers. Showed no sensitization and edema on skin after 72 hrs of application. The results of present study revealed that the prepared medicated sticks of *Abrus Precatorius* are convenient, equally effective, without any contamination chances on application and free from skin irritation.

KEYWORDS: *Abrus Precatorius*, medicated sticks.

INTRODUCTION

Candidiasis: Candidosis (Candidiasis, moniliasis) is an infection of the skin, mucosa and rarely of the internal organs, caused by yeast like fungus *Candida albicans*, normally present in vagina. *Candida albicans* is an ovoid or spherical budding cell, which produces pseudo-mycelia both in culture and in tissues. *Candida* species are normal inhabitants of the skin and mucosa. Over the last two decades, there has been a dramatic increase in the rate of superficial and invasive fungal infections. Approximately three-quarters of all women experience at least one episode of vulva-vaginal candidiasis during their life time and nearly half of them suffer from multiple episode. The manifestations of vulva vaginal candidiasis are often painful and uncomfortable and can include intense itching, irritation, vaginal discharge and dysuria. Gynecological skin infections are very common in most of the woman population especially in rural areas due to unhygienic maintenance. Many patients express difficulty in application of ointments, creams, gels etc. results in non-compliance and ineffective therapy. Recent advance in novel drug delivery systems (NDDS) aim to enhance safety and efficacy of drug molecules by formulating a convenient dosage form for application and to achieve better patient compliance. One such approach is medicated sticks.^[1-2] An advantage of this drug delivery system includes patient compliance; convenience and comfort ness for efficient treatment include application without fingertip, immediate onset of action, reduced dosage regimen and economy. *Abrus Precatorius*^[3-5] a herbal medication has anti bacterial and anti fungal activity commonly used in the treatment of several skin disorders as it is not available in such dosage form.^[6-7] *Abrus precatorius* is a slender, perennial climber that twines around trees, shrubs, and hedges. It has no special organs of attachment. It has a slender branch and a

cylindrical wrinkled stem with a smooth-textured brown bark. Leaves are glabrous with long internodes. Leaves alternate petiole, 5-13 cm (2-5 in) long, even-pinnately compound with 5-15 pairs of leaflets, these oval to oblong, to 1.8 cm (< 1 in) long, with margins entire. Flowers are small and pale violet in color with a short stalk, arranged in clusters. The ovary has a marginal placentation. The fruit, which is a pod, is flat, oblong and truncate- shaped with a sharp deflexed beak is about 3 to 4.5 cm long, 1.2 cm wide, and silky-textured. Each fruit contains from 3 to 5 oval-shaped seeds, about 0.6 cm. Cytotoxic and anthelmintic (seed); antiestrogenic (root); stimulates the cardiovascular system (aerial part); Abortifacient (seed and root), anodyne, aphrodisiac, antimicrobial, diuretic, emetic, expectorant, febrifuge, hemostat, laxative, purgative, refrigerant, sedative, vermin fuge, anti-fertility activity and antitumor activity. Objective of the present work was to develop such a NDDS of prepare Abrus Precatorius medicated derma sticks by heating and congealing method .for the treatment of skin infections.

MATERIALS AND METHODS

Abrus Precatorius was gift sample from Leads Pharmaceuticals., Hyderabad., T.S.. Stearyl alcohol pure, white petrolatum (Loba chemie Pvt. Ltd., Mumbai), Sodium lauryl sulphate, Cetyl Alcohol (S.D. fine chemicals ltd. Mumbai), Propylene glycol (Ranbaxy lab. Ltd., SAS Nagar), Methanol (Qualigens Fine Chemicals, Mumbai) were used.

Preparation of medicated derma sticks of Abrus Precatorius

Medicated derma sticks were prepared by heating and congealing according to the formulae (Table 1). Depending upon the weight, thickness and length of medicated derma sticks, the formulae was chosen for the incorporation of the drug. Stearyl alcohol^[8]/ Cetyl alcohol^[9] and white petroleum were melted in a china dish and heated this mixture up to 70°C. Dissolve sodium lauryl sulfate, propylene glycol in purified water and heat the solution to 70°C separately. Add the oleaginous phase slowly to the aqueous phase, stirring constantly and then the drug was added slowly with continuous stirring in order to get a uniform mixture in optimized formulation. The hot mixture was poured into the glass mould and cooled to get the desired shape of sticks. The stick was removed from the mould after 24 hours with the help of plunger and inserted into the medicated derma stick container (Table 1).

Evaluation of prepared medicated derma sticks

Three sticks were selected randomly and weighed individually. The individual weights were compared with the average weight for determination of weight variation. As the shape of the

stick is cylindrical the thickness and length was determined with the help of screw gauge and vernier callipers respectively. The average thickness was measured, by observing thickness at three different parts of the stick. (Table 2)

Anti microbial studies of prepared formulations^[10]

The anti microbial studies were carried out for the prepared formulations by cup-plate method using *Candida Albicans* as test organism. The cultures of *Candida albicans* were cultivated on Sabouraud's dextrose agar maintained on slants in the refrigerator ($4\pm 2^\circ\text{C}$). (Table 4)

Cup-plate method

The composition of Sabouraud's dextrose agar was taken in a 250 ml of conical flask and was dissolved in 100ml of distilled water. The pH was adjusted to 5.6. The medium was sterilized in an autoclave at 15 lbs for 20 minutes. After the completion of sterilization, the medium was kept aside at room temperature. 0.5 ml diluted suspension culture in NaCl 0.9% were added to 100 ml of medium at $47\pm 2^\circ\text{C}$ and used as inoculated layer. The medium (20 ml) was poured into a sterilized petridish to give a depth of 3-4 mm, and was assured that the layer of medium is uniform in thickness by placing petridish on a leveled surface. After solidifying the medium at room temperature, with the help of a sterile cork borer, cups of each 6 mm diameter were punched and scooped out from the petridish. Using sterile pipettes sample solutions (0.1 ml) of known concentration were fed into the cup. The petridish was then incubated for 24 hours at 37°C . After incubation the zone of inhibition was measured.

Preclinical studies: Primary skin irritation test in animals

This test is conducted on 3 healthy rabbits and guinea pigs (2 male and 1 female), which were fed with fresh food and water during the test period. 24 hours prior to test, the hair from the lower abdominal portion was shaved to expose sufficiently large test area. The test site was cleaned with surgical spirit then medicated stick is applied to test area. The test site was observed for erythematic and edema for 72 hrs. after application. This test was conducted to evaluate the irritancy of the prepared medicated stick on the intact skin of rabbits and guinea pigs. (Table 5,6)

Preclinical studies: Primary skin irritation test in healthy human being volunteers

3 Healthy Human Volunteers were selected for the study for each formulation. The test site was cleaned with surgical spirit then medicated stick is applied to test area. The test site was

observed for erythema and edema for 24 hrs. 48 hrs. & 72 hrs after application. This test was conducted to evaluate the irritancy of the prepared medicated stick on the intact skin. None of the prepared medicated sticks showed any erythematic or edema, indicating that the prepared formulations were non-irritant on the skin. . These studies were carried out under the guidance of qualified dermatologists with the permission of ethical committee of M. R. Medical College, Gulbarga.(Table 7)

Stability Studies

Short-term stability studies on the promising formulation MP-1 were carried out by storing the sticks at $27 \pm 2^\circ\text{C}$ for a period of three weeks. At intervals of one week the sticks were visually examined for any physical changes. (Table 3).

RESULTS AND DISCUSION

Steryl alcohol and Cetyl alcohol as stiffening agent while petrolatum used as emollient, propylene glycol and sodium lauryl sulphate were used as humectants and emulsifying agent respectively. Medicated sticks of abrus precatorius were prepared by heating and congealing method.. The sticks obtained were of uniform length, thickness and weight respectively. Antimicrobial studies revealed that the drug in formulation show equal zone of inhibition like pure drug. The preclinical studies in animals and healthy human volunteer revealed that the prepared formulations will be safe to use for topical applications (Fig 1 to4).

Table No. 1: Formula of medicated derma stick

Sl. No.	Ingredients	Quantity in gms.
1.	Abrus Precatorius powder	1.00
2.	Stearyl alcohol	15.00
3.	White petrolatum	20.00
4.	White Beeswax	5.00
5.	Sodium lauryl sulfate	1.50
6.	Propylene glycol	12.50
7.	Purified water (Q.S.)	100.00

Table No: 2 Characterization of prepared medicated derma sticks

Formulation code	Medicated Stick		
	Weight (gm)	Thickness (mm)	Length (cm)
	Mean $\pm$ SD	Mean $\pm$ SD	Mean $\pm$ SD
Abrus Precatorius	1.61 ± 0.02	6.62 ± 0.02	4.0 ± 0.01
Abrus Precatorius	1.61 ± 0.02	6.62 ± 0.02	4.0 ± 0.01
Abrus Precatorius	1.61 ± 0.02	6.62 ± 0.02	4.0 ± 0.01

Each reading is a mean of three determinations

Stability studies

Formulations containing were found to be stable at the temperatures and parameters tested.

Table No. 3: stability studies of *Abrus precatorius* derma sticks

Storage temperature	Time of analysis (days)	Physical appearance	pH
At room temperature 27 ± 2°C	15	No change	7.0
	30	No change	7.0
	45	No change	7.0
	60	No change	7.0
	75	No change	7.0
	90	No change	7.0
	105	No change	7.0
	120	No change	7.0
	135	No change	7.0
	150	No change	7.0
	165	No change	7.0
	180	No change	7.0

Each reading is a mean of three determinations



Pure drug




Plate-1



Pure drug



Plate- 2



Pure drug



Plate- 3

Figure 2: ANTIMICROBIAL STUDIES SHOWING THE COMPARATIVE ZONE OF INHIBITION OF DRUG AS PURE AND IN FORMULATION ABRUS PRECATORIUS

Table 4: ANTIMICROBIAL STUDIES SHOWING THE COMPARATIVE ZONE OF INHIBITION OF DRUG AS PURE AND IN FORMULATION ABRUS PRECATORIUS

Formulation code	Statistical zone of inhibition (mm) after 36 hrs.			Mean $\pm$ S.D.
	Zone 1	Zone 2	Zone 3	
Pure Drug	13	14	15	13.66 $\pm$ 0.57
Abrus Precatorius (Plate-1)	14	11	12	12.33 $\pm$ 1.53
Abrus Precatorius (Plate-2)	11	10	13	11.33 $\pm$ 1.53
Abrus Precatorius (Plate-3)	12	13	11	10.0 $\pm$ 1.00

Skin Irritation: All the formulation bases were subjected to skin irritation test in rabbits and guinea pigs. For three days of application.




Figure 3: Primary Skin irritation test of one of the group of Rabbits

Table No: 5 Skin irritation test data of prepared derma stick bases for Rabbits

Formulation code	Rabbits	Before application			After 24 hrs. of application			After 48 hrs. of application			After 72 hrs. of application		
		I	R	E	I	R	E	I	R	E	I	R	E
Abrus Precatorius	Male-I	x	x	x	x	x	x	x	x	x	x	x	x
	Male-II	x	x	x	x	x	x	x	x	x	x	x	x
	Female	x	x	x	x	x	x	x	x	x	x	x	x

I-Skin irritation,

R-Redness,

E-Erythema




Figure 4: Primary Skin irritation test of one of the group of Guinea Pigs


Table No: 6 Skin irritation test data of prepared derma stick bases for Guinea pigs

Formulation code	Guinea pigs	Before application			After 24 hrs. of application			After 48 hrs. of application			After 72 hrs. of application		
		I	R	E	I	R	E	I	R	E	I	R	E
Abrus Precatorius	Male-I	x	x	x	x	x	x	x	x	x	x	x	x
	Male-II	x	x	x	x	x	x	x	x	x	x	x	x
	Female	x	x	x	x	x	x	x	x	x	x	x	x

I-Skin irritation

R-Redness

E-Erythema



Group 1: (Abrus Precatorius)
Figure: 5 Primary Skin Irritation Test of Healthy Human Volunteers

Table No: 7 Skin irritation test data of prepared derma stick bases for Healthy Human Volunteers

Formula tion code	Human Volunteers	Before application			After 24 hrs. of application			After 48 hrs. of application			After 72 hrs. of application		
		I	R	E	I	R	E	I	R	E	I	R	E
Abrus Precatori us	Male-I	x	x	x	x	x	x	x	x	x	x	x	x
	Male-II	x	x	x	x	x	x	x	x	x	x	x	x
	Female	x	x	x	x	x	x	x	x	x	x	x	x

I-Skin irritation

R-Redness

E-Erythema

CONCLUSIONS

The present work is a unique piece of contribution to the drug industry. The results will be useful to industry R&D for further investigations.

ACKNOWLEDGEMENTS

We are very much thankful to M/S Leads Pharmaceuticals Pvt Ltd., Hyderabad., T.S. for the encouragement during the study period.

REFERENCES

1. Fuchs P, Schopflin G. Medicated sticks, United States patent, 1974; 3,856,931. Berlin.
2. Fuchs P, Schopflin G. Medicated sticks, United States patent, 3,211,618,931: Berlin: 1974.
3. Indian pharmacopoeia, Vol. II, 4th ed. New Delhi: The control of publications; 1996; 673.
4. British pharmacopoeia, Her Majesty's Stationary Office for the Department of Health; 3rd ed., London, 2008; II: 1920-22.
5. United States Pharmacopoeia, Vol. III, Port city press: Asian ed. US, 2007; 3154-55.
6. Kumar V, Cotran RS, Robbin SL. Basic pathology, 6th ed. Harcourt India pvt. Ltd: New Delhi, 2001; 705-7.
7. Berkow R. editor. The merck manual, 14th ed. Merck Sharp and Dohme: Merck Co. Inc, 1982; 2046.
8. Aniley W, Paul WJ. Hand book of pharamaceutical excipients: profile of stearyl alcohol. The pharamaceutical Press: London, 1994; 498.
9. Aniley W, Paul WJ. Hand book of pharamaceutical excipients: profile of cetyl alcohol. The pharamaceutical Press: London, 1994; 99.

10. Indian pharmacopoeia, Vol. II, 4th ed. New Delhi: The control of publications, 1996; A-147.
11. Bango R, Jayakar B. Diffusion studies on salicylic acid ointment through rabbit skin. The Indian Journal of Hospital Pharmacy; 1997; 51- 2.