

2-METHYL IMIDAZOLIUMBROMO CHROMATE: A NEW AND EFFICIENT REAGENT FOR OXIDATION OF ALIPHATIC ALCOHOLIC COMPOUNDS

Kyadare G. N.*

Department of chemistry, Degloor College, Degloor Di. Nanded- 431717, S. R. T. M.

University Nanded (MS).

Article Received on
04 April 2023,

Revised on 25 April 2023,
Accepted on 15 May 2023

DOI: 10.20959/wjpr20238-28164

***Corresponding Author**

Kyadare G. N.

Department of chemistry,
Degloor College, Degloor
Di. Nanded- 431717, S. R.
T. M. University Nanded
(MS).

ABSTRACT

The new chromium (VI) oxidizing reagent 2-methyl imidazoliumbromochromate (IBC) was prepared and characterized. The 2- methyl IBC has been found to be stable and an efficient solid reagent which can be easily prepared in good yield. It act as an efficient brominating agent for hydroxylated aromatic compounds as well as good oxidizing agent for the conversion of alcohols to carbonyl compounds with good yield.

KEYWORDS: 2-methyl Imidazoliumbromochromate, Oxidation: Alcohols.

INTRODUCTION

Chromium (VI) complexes are among the most interesting transition metal complexes that have great attention over the past decades. Due to their appealing catalytic and oxidizing properties, they have been widely explored since the very beginning of organic chemistry. The topic remains of current interest as exemplified by the huge number of reactions involving the use of chromium (VI) reagents. Moreover, a plethora of chromium reagents and procedures have been proposed and described extensively. The main objective of present study is to synthesis of new chromium (VI) reagent for the oxidation of different organic substrates and encourages its various contributions to the important and exciting field of synthetic organic chemistry.^[2-3] The oxidation of alcohols is the important basic reaction in organic chemistry and most commonly used method for the synthesis of

aldehydes and ketones. Therefore, the search for new oxidizing agent is of interest to synthetic organic chemists.

In recent years some chromium (VI) reagents for the effective and selective oxidation of alcohols under mild conditions have been developed with some success.^[4] Some of reagents like pyridinium dichromate^[5], pyridinium fluorochromate^[6], quinolinium chlorochromate^[7], quinolinium fluorochromate^[8], imidazolium chlorochromate^[9], prolinium chlorochromate^[10], caffelinillium chlorochromate^[11], are reported as oxidizing agents while some reagents like pyridinium bromochromate^[12], benzimidazolium bromochromate^[13], quinoxalinium bromochromate^[14] works as oxidizing as well as good brominating agent.

Oxidation states of chromium metal includes Cr+2, Cr+3, Cr+6 which are stable states whereas Cr+4 and Cr+5 are unstable in water and they form Cr+3 and Cr+6. Cr+6 is used as oxidant for various reactions. Chromium (VI) exists as acetochromate ion in presence of 97% acetic acid¹⁵. New chromium species are formed on adding anions like Cl⁻, Br⁻, F⁻, and So4²⁻ anions. The chemistry of chromium (VI) complexes has advantage in both synthetic organic chemistry and medicinal field because of its high reactivity. Nowadays, the development of new and versatile chromium (VI) reagents¹⁶⁻²¹ for oxidizing organic molecules is of great interests and studied extensively. Oxides and oxyacid's of Cr (VI) are powerful oxidants and chromic esters have been proposed as intermediates in Cr(VI) oxidations.


Experimental

General procedure for oxidation of alcohols



To a solution of substrate (0.01 mol) in acetic acid (15ml) 2M-Imidazolium bromochromate (0.278g, 1mmol) was added and the mixture was stirred magnetically for 2-3 h. The progress of reaction was monitored by TLC. After completion of reaction, reaction mixture was distilled by circulation of ice cold water through condenser, extracted with ether and filtered through silica column. Finally it was washed with ether. On evaporation of ether product is obtained.^[22]

Probable reaction mechanism of oxidation of alcohols

Reaction




Mechanism



Preparation of 2-methyl Imidazoliumbromochromate

This was synthesized^[6] by taking 20g (0.2 mol) of chromium trioxide in water (25ml), cooled to 0°C. To this solution, 47% HBr (38ml, 0.21mol) was added slowly with constant stirring. The content were cooled to 0°C. and then 2-methyl Imidazole (17gm 0.2 mol) was added over 15-20 min. to give a brown solid. The reaction mixture was chilled for 4-5 hr. The dark brown crystals were then filtered and dried. The product was recrystallized from aqueous acetic acid (40:60 v/v) its purity was checked by TLC and confirmed by melting point and mass analysis.


Structure of 2-methyl bromochromate



2methyl imidazolium bromochromate

Oxidation of alcohols by 2-methyl IBC: A study of oxidative behavior of 2-methyl IBC with primary and secondary alcohol in acetic acid was undertaken (Scheme 1) and the results are presented in table 1. The results of oxidation obtained with 2M-IBC are satisfactory in comparison with other reagent considering the amount of oxidant, acidity, percentage yield and reaction time, respectively.

Reaction



Scheme-1

Table 1

Oxidation of alcohols by 2M-Imidazolium bromochromate

Sr. No.	Starting material	Product	B.P.(0C) (Lit)
1	n-Hexyl alcohol	1-Hexanal	129 (130) [23]
2	Isobutyl alcohol	2-Butanone	79 (82) [23]
3	n-Butyl alcohol	1-Butanal	74 (75) [23]
4	Benzyl alcohol	Benzaldehyde	180 (179) [23]
5	n-Propyl alcohol	Propanal	50 (49) [23]
6	Isopropyl alcohol	Acetone	55 (56) [23]
7	Amyl alcohol	1-Pentanal	105 (104) [23]
8	Isoamyl alcohol	2-Pentanone	101 (102) [23]

REFERENCES

1. Shankar A.K., Carlos Cervantes, Herminia Loza-Tavera and Avudainayagam S., Chromium toxicity in plants, *Environment International*, 2005; 31: 739–753.
2. A. Butler, J.V. Walker, *Chem. Rev.*, 1993; 93: 1937.
3. L.F. Fieser, M. Fieser, *Reagent for Organic Synthesis*, Wiley, New York, 1967; 1–11.
4. E.J. Corey, G. Schmidt, *Tetrahedron Lett.*, 1979; 399.
5. M.N. Bhattacharjee, M.K. Chaudhari, H.S. Dasgupta, N. Roy, *Synthesis*, 1982; 588.
6. J. Singh, P.S. Kalsi, G.S. Jawanda, B.R. Chhabra, *Chem. Ind.*, 1986; 751.
7. V. Murugesan, A. Pandurangan, *Indian J. Chem.*, 1992; 31B: 377.
8. K. Sunggak, C. Dong, *Bull. Chem. Soc. Jpn.*, 1986; 59: 3297.

9. M. Mamaghani, F. Shirini, F. Parsa, Russ. J. Org. Chem., 2002; 38(8): 1113.
10. F. Shirini, I. Mohammadpoor-Baltork, Z. Hejazi, P. Heravi, Bull. Korean Chem. Soc., 1999; 29(5): 763.
11. N. Narayana, T.R. Balasubramanian, Indian J. Chem., 1986; 25B: 228.
12. B. Ozgun, N. Degimenbasi, Synth. Commun, 1990; 20(21): 3359.
13. B. Ozgun, N. Degimenbasi, G. U. J. Sci., 2006; 19(1): 9.
14. Chaudhuri M.K., Chettri S.K., Lyndem S., Paul P.C. and Srinivas P., Quinolinium Fluorochromate (QFC), C₉H₇NH[CrO₃F]: An Improved Cr(VI)-Oxidant for Organic Substrates, Bull. Chem. Soc. Jpn., 1994; 67: 1894-1898.
15. Bhattacharjee M.N., Choudhari M.K., Dasgupta H.S., Roy N. and Khating D.T., Pyridinium Fluorochromate: A New and Efficient Oxidant for Organic Synthesis, Synthesis, 1982; 1982(07): 588- 590.
16. Corey E.J., Barette E.P.M. and Margrious P.A., A new Cr(VI) reagent for the catalytic oxidation of secondary alcohols to ketones, Tetrahedron Lett., 1985; 26(48): 5855-5858.
17. Climinale F., Camporeale M., Mello R., Troisi L. and Curci R., Oxidation of Tertiary Amines by Chromium(VI) oxide diperoxide, J. Chem. Soc., Perkin Trans, 1989; 2: 417-423.
18. Sharma G.G. and Mahanti M.K., Bull. Soc. Chem. Fr., 1991; 128: 449.
19. Balasubramanian K. and Pratibha V., Indian J. Chem. Sec B, 1986; 25: 326.
20. Narayana B. and Cherian T., Rapid Spectrophotometric Determination of Trace Amounts of Chromium using Variamine Blue as a Chromogenic Reagent, J. Braz. Chem. Soc., 2005; 16(2): 197-201.
21. Y. Martinez, M.A. DelasHeras, J.J. Vaquero, J.L. Garcianario, J. Alvarezbuilla, Tetrahedron, 1995; 31: 8513.
22. S.V. Khansole et al. / Chinese Chemical Letters, 2009; 20: 256–260 259.
23. Vogel's Text book of Organic Chemistry, 4th ed., Longman Publisher, England, 1978.